



Alliance Pediatrics

Antihistamine Only	Benadryl Allergy Liquid 12.5 mg/ 5ml	Benadryl I Dye-free Allergy Liquid 12.5 mg/ 5ml	Benadryl Allergy FASTMELT® Tablets 12.5 mg/ tab
Flavor	Cherry	Bubble Gum	Grape
Dosing Frequency	Every 4-6 hours	Every 4-6 hours	Every 4-6 hours
*5 lbs. 8 oz. – 7 lbs. 10 oz.	1.5 ml every 6 hours as needed	1.5 ml every 6 hours as needed	N/A
*7 lbs. 11 oz. – 9 lbs. 14 oz.	2 ml every 6 hours as needed	2 ml every 6 hours as needed	N/A
*9 lbs. 15 oz. – 12 lbs. 1 oz.	3 ml every 6 hours as needed	3 ml every 6 hours as needed	N/A
*12 lbs. 2 oz. – 14 lbs. 4 oz.	3 ml every 6 hours as needed	3 ml every 6 hours as needed	N/A
*14 lbs. 5 oz. – 16 lbs. 7 oz.	3.5 ml every 6 hours as needed	3.5 ml every 6 hours as needed	N/A
*16 lbs. 8 oz. – 18 lbs. 10 oz.	4 ml every 6 hours as needed	4 ml every 6 hours as needed	N/A
*18 lbs. 11 oz. – 20 lbs. 14 oz.	4.5 ml every 6 hours as needed	4.5 ml every 6 hours as needed	N/A
*20 lbs. 15 oz. – 23 lbs. 15 oz.	5 ml every 6 hours as needed OR one 12.5 mg tablet	5 ml every 6 hours as needed OR one 12.5 mg tablet	One 12.5 mg tablet
24 lbs. – 28 lbs. 6 oz.	6 ml every 6 hours as needed	6 ml every 6 hours as needed	One 12.5 mg tablet
28 lbs. 7 oz. – 32 lbs. 12 oz.	7 ml every 6 hours as needed	7 ml every 6 hours as needed	One 12.5 mg tablet
32 lbs. 13 oz. – 37 lbs. 2 oz.	8 ml every 6 hours as needed	8 ml every 6 hours as needed	One 12.5 mg tablet
37 lbs. 6 oz. – 41 lbs. 9 oz.	9 ml every 6 hours as needed	9 ml every 6 hours as needed	One 12.5 mg tablet
42 lbs. and up	10 ml every 6 hours as needed OR two 12.5 mg tablets	10 ml every 6 hours as needed OR two 12.5 mg tablets	10 ml every 6 hours as needed OR two 12.5 mg tablets
6-8 yrs. 48-59 lbs.	1 to 2 tsp.	1 to 2 tsp.	One or Two 12.5 mg tablets
9-10 yrs. 60-71 lbs.	1 to 2 tsp.	1 to 2 tsp.	One or Two 12.5 mg tablets
11 yrs. 71-95 lbs.	1 to 2 tsp.	1 to 2 tsp.	One or Two 12.5 mg tablets
12+ yrs. 96+ lbs.	2 tsp.	2 tsp.	One or Two 12.5 mg tablets

*Please consult the office prior to using this medication as there are risk of overdose, especially when used with other medicines. This chart is a guide to dosing and each child may have qualities that make this dose of even this medication inappropriate.